



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2011

1,3-Bis(pyridin-2-yl)-1H-benzimidazol-3-ium tetrafluoridoborate

Grieco, G ; Blacque, O ; Berke, H

Abstract: The asymmetric unit of the title compound, C(17)H(13)N(4)(+)center dot BF(4) , contains one half of the benzimidazolium cation and one half of the tetrafluoridoborate anion, with crystallographic mirror planes bisecting the molecules. One F atom of the tetrafluoridoborate is equally disordered about a crystallographic mirror plane. In the crystal, C-H center dot center dot center dot F interactions link the cations and anions into layers parallel to (100). The crystal packing is further stabilized by F center dot center dot center dot pi contacts involving the tetrafluoridoborate anions and the five-membered rings [F center dot center dot center dot centroid = 2.811 (2) angstrom].

DOI: <https://doi.org/10.1107/S1600536811027942>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-60388>

Journal Article

Published Version

Originally published at:

Grieco, G; Blacque, O; Berke, H (2011). 1,3-Bis(pyridin-2-yl)-1H-benzimidazol-3-ium tetrafluoridoborate. Acta Crystallographica. Section E, Structure Reports Online, 67(Part 8):O2066-U805.

DOI: <https://doi.org/10.1107/S1600536811027942>

1,3-Bis(pyridin-2-yl)-1*H*-benzimidazol-3-ium tetrafluoridoborate

Gabriele Grieco, Olivier Blacque* and Heinz Berke

Institute of Inorganic Chemistry, University of Zürich, Winterthurerstrasse 190, 8057 Zürich, Switzerland

Correspondence e-mail: oblacque@aci.uzh.ch


Received 29 June 2011; accepted 12 July 2011

Key indicators: single-crystal X-ray study; $T = 183$ K; mean $\sigma(\text{C}—\text{C}) = 0.002$ Å; disorder in solvent or counterion; R factor = 0.042; wR factor = 0.117; data-to-parameter ratio = 15.5.

The asymmetric unit of the title compound, $\text{C}_{17}\text{H}_{13}\text{N}_4^+\cdot\text{BF}_4^-$, contains one half of the benzimidazolium cation and one half of the tetrafluoridoborate anion, with crystallographic mirror planes bisecting the molecules. One F atom of the tetrafluoridoborate is equally disordered about a crystallographic mirror plane. In the crystal, $\text{C}—\text{H}\cdots\text{F}$ interactions link the cations and anions into layers parallel to (100). The crystal packing is further stabilized by $\text{F}\cdots\pi$ contacts involving the tetrafluoridoborate anions and the five-membered rings [$\text{F}\cdots\text{centroid} = 2.811$ (2) Å].

Related literature

For applications of N,N' -bis(2-pyridyl)aryldiamines, see: Stoessel *et al.* (2010); Goldfarb (2009) and of imidazolium salts, see: Berlin *et al.* (2007); Bold *et al.* (2005); Huang *et al.* (2005); Murakami *et al.* (2007); Teles *et al.* (1996). For pharmaceuticals based on the aniline–pyridine scaffold, see: Kim *et al.* (1996); Wu *et al.* (2001). For the synthesis of the starting material N,N' -bis(pyridin-2-yl)benzene-1,2-diamine, see: Gdaniec *et al.* (2004).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{13}\text{N}_4^+\cdot\text{BF}_4^-$

$M_r = 360.12$

Orthorhombic, $Pnma$

$a = 7.3412$ (2) Å

$b = 17.5051$ (5) Å

$c = 12.2426$ (3) Å

$V = 1573.28$ (7) Å³

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 0.13$ mm^{−1}

$T = 183$ K

$0.44 \times 0.31 \times 0.11$ mm

Data collection

Oxford Diffraction Xcalibur diffractometer with a Ruby detector

Absorption correction: analytical

[*CrysAlis PRO* (Oxford

Diffraction, 2010) based on Clark

& Reid (1995)]

$T_{\min} = 0.964$, $T_{\max} = 0.991$

8816 measured reflections

2014 independent reflections

1543 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.022$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$

$wR(F^2) = 0.117$

$S = 1.08$

2014 reflections

130 parameters

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\max} = 0.27$ e Å^{−3}

$\Delta\rho_{\min} = -0.36$ e Å^{−3}

Table 1

Hydrogen-bond geometry (Å, °).

$D—H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
$\text{C1}—\text{H1}\cdots\text{F3}$	0.95 (2)	2.14 (2)	3.094 (2)	178 (2)
$\text{C9}—\text{H9}\cdots\text{F1}^1$	0.93	2.62	3.4759 (19)	154

Symmetry code: (i) $-x + \frac{3}{2}, -y, z - \frac{1}{2}$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *Mercury* (Macrae *et al.*, 2006), *ORTEP-3 for Windows* (Farrugia, 1997) and *POV-RAY for Windows* (Cason, 2003); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *publCIF* (Westrip, 2010).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2291).

References

- Berlin, J. M., Campbell, K., Ritter, T., Funk, T. W., Chlenov, A. & Grubbs, R. H. (2007). *Org. Lett.* **9**, 1339–1342.
- Bold, M., Lennartz, C., Prinz, M., Schmidt, H.-W., Thelakkat, M., Baete, M., Neuber, C., Kowalsky, W., Schildknecht, C. & Johannes, H.-H. (2005). PCT Int. Appl. WO 2005019373.
- Cason, C. J. (2003). *POV-RAY*. Persistence of Vision Raytracer Pty. Ltd, Victoria, Australia.
- Clark, R. C. & Reid, J. S. (1995). *Acta Cryst.* **A51**, 887–897.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Gdaniec, M., Bensemann, I. & Połowski, T. (2004). *Acta Cryst.* **C60**, o215–o216.
- Goldfarb, D. S. (2009). US Patent Appl. Publ. US 20090163545.
- Huang, W., Guo, J., Xiao, Y., Zhu, M., Zou, G. & Tang, J. (2005). *Tetrahedron*, **61**, 9783–9790.
- Kim, H.-J., Han, Y.-H., Chung, S.-J., Lee, M.-H. & Shim, C.-K. (1996). *Arch. Pharm. Res.* **19**, 297–301.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.

- Murakami, T., Yagi, K., Ichijima, S., Igarashi, T. & Satou, T. (2007). WO 2007034985.
- Oxford Diffraction (2010). *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, England.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Stoessel, P., Heil, H., Joosten, D., Pflumm, C. & Gerhard, A. (2010). PCT Int. Appl. WO 2010099852.
- Teles, J. H., Melder, J.-P., Ebel, K., Schneider, R., Gehrer, E., Harder, W., Brode, S., Enders, D., Breuer, K. & Raabe, G. (1996). *Helv. Chim. Acta*, **79**, 61–83.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Wu, S. N., Jan, C. R. & Chiang, H. T. (2001). *J. Investig. Med.* **49**, 522–533.

supplementary materials

Acta Cryst. (2011). E67, o2066–o2067 [doi:10.1107/S1600536811027942]

1,3-Bis(pyridin-2-yl)-1*H*-benzimidazol-3-ium tetrafluoroborate

G. Grieco, O. Blacque and H. Berke

Comment

N,N'-bis(2-pyridyl)aryldiamines are an important class of compounds that are useful as intermediates in the syntheses of organic electroluminescent devices (LED) (Stoessel *et al.*, 2010), or as compounds useful for altering the lifespan of eukaryotic organisms in the yeast (Goldfarb, 2009). Imidazolium salts based on the benzoimidazole scaffold are used in electronics (Bold *et al.*, 2005; Murakami *et al.*, 2007) and in catalytic processes involving the use of the benzimidazolium salt as it is (Teles *et al.*, 1996), or bound to a metal (Berlin *et al.*, 2007; Huang *et al.*, 2005).

We used the *N,N'*-bis(2-pyridyl)-benzene-1,2-diamine compound as a starting material for the synthesis of the title compound, a new imidazolium salt. Despite the fact that the synthesis of the starting material *N,N'*-bis(pyridin-2-yl)benzene-1,2-diamine had been reported previously (Gdaniec *et al.*, 2004), we optimized the procedure to obtain a better yield (91 instead of 70%) using the reaction reported in the Experimental section (Fig. 4) with a microwave technique. A similar procedure was successfully used to synthesize the title compound in a very high yield (99.7%). It is the first example of a coupling between a halo pyridine and an aniline made with microwave heating that does not imply the use of any metal. This method to couple halo pyridines and anilines can be very useful to produce compounds with pharmaceutical activity since many pharmaceuticals are based on the aniline-pyridine scaffold (Kim *et al.*, 1996; Wu *et al.*, 2001).

The asymmetric unit of the title compound, $C_{17}H_{13}N_4^+.BF_4^-$, contains one half-molecule of the benzimidazolium cation and one half-molecule of the anion, crystallographic mirror planes bisecting the molecules (Fig. 1). One F atom of the tetrafluoroborate is disordered over two positions. The second position being generated by a crystallographic mirror plane, the site-occupancy factor is 0.5. The benzimidazole and pyridine rings are not coplanar, the dihedral angle between the mean planes is 26.67 (4)°.

In the crystal intermolecular C—H...F interactions link the cations and anions into layers parallel to the (100) crystallographic plane (Fig. 2, Table 1). The crystal packing is further stabilized by F... π contacts (Fig. 3) involving the tetrafluoroborate anions and the five-membered rings of the benzimidazole rings [$F\cdots\text{centroid} = 2.811$ (2) Å].

Experimental

To benzene-1,2-diamine (2.7 g, 24.97 mmol) in a microwave vial, 2-chloropyridine (9 ml, 46.95 mmol) was added. The vial was then closed with a cap consisting of a Teflon septum and the reaction mixture was heated for 35 mins at 458 K. Monitoring with TLC (thin layer chromatography) and GC—MS (gas chromatography - mass spectrometry) showed that the benzene-1,2-diamine was consumed after this time and the mixture was allowed to cool to room temperature. The crude mixture was dissolved in water (15 ml) and dropped into a solution of concentrated ammonia in water (25 ml of NH_4OH 24.5% in 250 ml of water). The resulting pink-red precipitate was filtered off, washed with water (50 ml) and was subsequently dried in air to give *N,N'*-bis(pyridin-2-yl)benzene-1,2-diamine. Further recrystallization from ethanol gave a very pure product (yield: 2.902 g, 91%).

To *N,N'*-bis(pyridin-2-yl)benzene-1,2-diamine (500 mg, 1.91 mmol) in a microwave vial, finely ground ammonium tetrafluoroborate (204 mg, 1.91 mmol) was added followed by the triethyl orthoformate (10 ml, 59 mmol). The vial with the red pink suspension was then closed with a cap consisting of a Teflon septum and the reaction mixture was heated for 25 minutes at 413 K and then for further 20 minutes to 433 K. After that time inside the vial a blue-violet solid was present and TLC analysis showed that the starting material had been consumed. The solid was separated off and then stirred with ethyl acetate (3x50 ml) for 15 minutes. It was then collected by suction filtration, washed with diethyl ether and dried to afford a deep violet-blue compound (yield: 686 mg, 99.7%). Recrystallization from a hot saturated solution of water/methanol (9/2) afforded red plate-like crystals of the title compound, suitable for X-ray analysis. Spectroscopic data for the title compound are given in the archived CIF.

Refinement

One F atom of the tetrafluoroborate ion is disordered over two positions around a mirror plane (site-occupancy factor of 1/2). H-atom H1 was located in a difference Fourier map and was freely refined. The C-bound H-atoms were included in calculated positions and treated as riding atoms: C—H = 0.93 Å with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of the title compound, with the numbering scheme and displacement ellipsoids drawn at the 50% probability level (H-atoms are shown as spheres of arbitrary size). Symmetry code: (i) $-x + 3/2, -y, z - 1/2$.




Fig. 2. View normal to (100) of the two-dimensional network of the title compound, formed by intermolecular C—H...F interactions (dashed lines; see Table 1 for details).




Fig. 3. View normal to (001) of the three-dimensional network of the title compound, formed by intermolecular B—F... π interactions (dashed lines; see Comment section for details).




Fig. 4. Reaction scheme.

1,3-Bis(pyridin-2-yl)-1*H*-benzimidazol-3-ium tetrafluoridoborate

Crystal data

$\text{C}_{17}\text{H}_{13}\text{N}_4^+ \cdot \text{BF}_4^-$

$M_r = 360.12$

Orthorhombic, *Pnma*

$F(000) = 736$

$D_x = 1.52 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Hall symbol: -P 2ac 2n	Cell parameters from 4054 reflections
$a = 7.3412$ (2) Å	$\theta = 2.8\text{--}32.6^\circ$
$b = 17.5051$ (5) Å	$\mu = 0.13\text{ mm}^{-1}$
$c = 12.2426$ (3) Å	$T = 183\text{ K}$
$V = 1573.28$ (7) Å ³	Plate, red
$Z = 4$	$0.44 \times 0.31 \times 0.11\text{ mm}$

Data collection

Oxford Diffraction Xcalibur diffractometer with a Ruby detector	1543 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.022$
Absorption correction: analytical [<i>CrysAlis PRO</i> (Oxford Diffraction, 2010) based on expressions derived by Clark & Reid (1995)]	$\theta_{\text{max}} = 28.3^\circ$, $\theta_{\text{min}} = 3.2^\circ$
$T_{\text{min}} = 0.964$, $T_{\text{max}} = 0.991$	$h = -9 \rightarrow 9$
8816 measured reflections	$k = -20 \rightarrow 23$
2014 independent reflections	$l = -12 \rightarrow 16$

Refinement

Refinement on F^2	0 restraints
Least-squares matrix: full	H atoms treated by a mixture of independent and constrained refinement
$R[F^2 > 2\sigma(F^2)] = 0.042$	$w = 1/[\sigma^2(F_o^2) + (0.062P)^2 + 0.2418P]$
$wR(F^2) = 0.117$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.08$	$(\Delta/\sigma)_{\text{max}} < 0.001$
2014 reflections	$\Delta\rho_{\text{max}} = 0.27\text{ e \AA}^{-3}$
130 parameters	$\Delta\rho_{\text{min}} = -0.36\text{ e \AA}^{-3}$

Special details

Experimental. Spectroscopic data for the title compound:

¹H-NMR in CD₃CN: $\delta = 10.07$ (s, 1H), $\delta = 8.837$ (dd, 2H), $\delta = 8.48$ (m, 2H), $\delta = 8.29$ (m, 2H), $\delta = 8.04$ (t, 1H), $\delta = 8.01$ (t, 1H), $\delta = 7.87$ (m, 2H) $\delta = 7.77$ (m, 2H).

¹³C-NMR in CD₃CN: $\delta = 150.95$, 141.60, 129.49, 126.93, 118.83, 118.26, 116.86.

¹⁹F-NMR in CD₃CN: $\delta = -152.32$.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -

supplementary materials

factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	0.7110 (2)	0.25	0.56060 (15)	0.0232 (4)	
C2	0.64692 (16)	0.21032 (7)	0.39260 (10)	0.0211 (3)	
C3	0.60860 (17)	0.16850 (8)	0.29898 (10)	0.0243 (3)	
H3	0.6088	0.1154	0.299	0.029*	
C4	0.57020 (17)	0.20992 (8)	0.20603 (11)	0.0263 (3)	
H4	0.5436	0.1841	0.1416	0.032*	
C5	0.69854 (17)	0.11073 (8)	0.53968 (11)	0.0245 (3)	
C6	0.6441 (2)	0.09450 (9)	0.64542 (12)	0.0336 (3)	
H6	0.6026	0.1324	0.6925	0.04*	
C7	0.6546 (2)	0.01864 (9)	0.67741 (13)	0.0397 (4)	
H7	0.622	0.0045	0.748	0.048*	
C8	0.7136 (2)	−0.03573 (9)	0.60414 (14)	0.0374 (4)	
H8	0.7192	−0.087	0.624	0.045*	
C9	0.7640 (2)	−0.01265 (8)	0.50081 (14)	0.0345 (4)	
H9	0.8038	−0.0496	0.4517	0.041*	
N1	0.68805 (14)	0.18768 (6)	0.49998 (9)	0.0223 (3)	
N2	0.75864 (17)	0.06028 (7)	0.46749 (10)	0.0302 (3)	
F1	0.58623 (16)	0.18658 (7)	0.89742 (12)	0.0763 (4)	
F2	0.8041 (3)	0.2711 (2)	0.98569 (15)	0.0680 (16)	0.5
F3	0.8073 (2)	0.25	0.80661 (11)	0.0577 (4)	
B1	0.6969 (3)	0.25	0.89769 (19)	0.0338 (5)	
H1	0.743 (3)	0.25	0.6360 (19)	0.024 (5)*	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0249 (9)	0.0230 (10)	0.0217 (9)	0	−0.0010 (7)	0
C2	0.0210 (6)	0.0224 (7)	0.0200 (6)	0.0008 (5)	0.0019 (5)	0.0021 (5)
C3	0.0254 (6)	0.0229 (7)	0.0245 (7)	−0.0016 (5)	0.0012 (5)	−0.0029 (5)
C4	0.0263 (6)	0.0313 (7)	0.0212 (6)	−0.0013 (5)	0.0002 (5)	−0.0043 (5)
C5	0.0257 (6)	0.0215 (7)	0.0263 (7)	−0.0014 (5)	−0.0046 (5)	0.0030 (5)
C6	0.0450 (8)	0.0289 (8)	0.0269 (7)	−0.0052 (6)	−0.0002 (6)	0.0010 (6)
C7	0.0521 (9)	0.0371 (9)	0.0297 (7)	−0.0092 (7)	−0.0045 (7)	0.0114 (7)
C8	0.0416 (9)	0.0249 (8)	0.0457 (9)	−0.0009 (6)	−0.0092 (7)	0.0117 (7)
C9	0.0388 (8)	0.0235 (8)	0.0411 (8)	0.0055 (6)	−0.0036 (6)	0.0020 (6)
N1	0.0258 (5)	0.0211 (6)	0.0200 (5)	0.0001 (4)	−0.0006 (4)	0.0015 (4)
N2	0.0353 (6)	0.0230 (6)	0.0325 (6)	0.0033 (5)	0.0002 (5)	0.0019 (5)
F1	0.0558 (7)	0.0436 (7)	0.1295 (12)	−0.0010 (5)	0.0195 (7)	0.0143 (7)
F2	0.0400 (9)	0.132 (5)	0.0315 (9)	−0.0042 (12)	−0.0083 (7)	−0.0174 (14)
F3	0.0646 (10)	0.0810 (12)	0.0276 (7)	0	0.0114 (6)	0
B1	0.0284 (11)	0.0494 (16)	0.0236 (11)	0	−0.0016 (9)	0

Geometric parameters (Å, °)

C1—N1	1.3302 (15)	C6—C7	1.387 (2)
C1—H1	0.95 (2)	C6—H6	0.93
C2—C3	1.3887 (18)	C7—C8	1.378 (2)
C2—C2 ⁱ	1.389 (3)	C7—H7	0.93
C2—N1	1.4060 (16)	C8—C9	1.379 (2)
C3—C4	1.3784 (18)	C8—H8	0.93
C3—H3	0.93	C9—N2	1.3408 (19)
C4—C4 ⁱ	1.403 (3)	C9—H9	0.93
C4—H4	0.93	F1—B1	1.3756 (18)
C5—N2	1.3249 (18)	F2—B1	1.384 (3)
C5—C6	1.384 (2)	F3—B1	1.379 (3)
C5—N1	1.4341 (17)		
N1—C1—N1 ⁱ	110.20 (16)	C8—C7—H7	120.2
N1—C1—H1	124.88 (9)	C6—C7—H7	120.2
N1 ⁱ —C1—H1	124.88 (9)	C7—C8—C9	118.64 (14)
C3—C2—C2 ⁱ	121.81 (8)	C7—C8—H8	120.7
C3—C2—N1	131.81 (12)	C9—C8—H8	120.7
C2 ⁱ —C2—N1	106.37 (7)	N2—C9—C8	123.39 (15)
C4—C3—C2	116.46 (13)	N2—C9—H9	118.3
C4—C3—H3	121.8	C8—C9—H9	118.3
C2—C3—H3	121.8	C1—N1—C2	108.53 (11)
C3—C4—C4 ⁱ	121.74 (8)	C1—N1—C5	125.06 (12)
C3—C4—H4	119.1	C2—N1—C5	126.39 (11)
C4 ⁱ —C4—H4	119.1	C5—N2—C9	116.19 (13)
N2—C5—C6	125.68 (13)	F1—B1—F1 ⁱ	107.61 (19)
N2—C5—N1	114.70 (12)	F1—B1—F3	110.21 (13)
C6—C5—N1	119.61 (12)	F1—B1—F2 ⁱ	97.01 (18)
C5—C6—C7	116.38 (14)	F1—B1—F2	123.6 (2)
C5—C6—H6	121.8	F1 ⁱ —B1—F2	97.01 (18)
C7—C6—H6	121.8	F3—B1—F2	107.17 (18)
C8—C7—C6	119.69 (14)		

Symmetry codes: (i) $x, -y+1/2, z$.

Hydrogen-bond geometry (Å, °)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C1—H1 \cdots F3	0.95 (2)	2.14 (2)	3.094 (2)	178.(2)
C9—H9 \cdots F1 ⁱⁱ	0.93	2.62	3.4759 (19)	154

Symmetry codes: (ii) $-x+3/2, -y, z-1/2$.

Fig. 1




Fig. 2




Fig. 3




Fig. 4

